




DON $\mu\text{g}/\text{kg}$ in UK Barley from crop years 2015 to 2019
(EU MRL was set at 1250ppb ($\mu\text{g}/\text{kg}$) for DON in cereals for processing in 2007)




3 & 15-AcDON $\mu\text{g}/\text{kg}$ in UK Barley from crop years 2015 to 2019




NIV $\mu\text{g}/\text{kg}$ in UK Barley from crop years 2015 to 2019




HT2 & T2 $\mu\text{g}/\text{kg}$ in UK Barley from crop years 2015 to 2019
(EU set a combined Guidance Level of 200 for unprocessed cereals in 2013)




Fusarenone-X $\mu\text{g}/\text{kg}$ in UK barley from 2015 to 2019 crop years




Diacetoxyscirpenol $\mu\text{g}/\text{kg}$ in UK barley from 2015 to 2019 crop years




Neosolaniol $\mu\text{g}/\text{kg}$ in UK barley from 2015 to 2019 crop years



Fusarenone-X $\mu\text{g}/\text{kg}$ in malt made from UK barley in crop years 2015 to 2019



Diacetoxyscirpenol $\mu\text{g}/\text{kg}$ in malt made from UK barley in crop years 2015 to 2019



Neosolaniol $\mu\text{g}/\text{kg}$ in malt made from UK barley in crop years 2015 to 2019

