




DON $\mu\text{g}/\text{kg}$ in UK Barley from crop years 2015 to 2019
(EU MRL was set at 1250ppb ($\mu\text{g}/\text{kg}$) for DON in cereals for processing in 2007)




3 & 15-AcDON $\mu\text{g}/\text{kg}$ in UK Barley from crop years 2015 to 2019




NIV $\mu\text{g}/\text{kg}$ in UK Barley from crop years 2015 to 2019




HT2 & T2 $\mu\text{g}/\text{kg}$ in UK Barley from crop years 2015 to 2019
(EU Guidance Level is currently set at 200ppb ($\mu\text{g}/\text{kg}$) in unprocessed cereals)




Fusarenone-X (FUS) $\mu\text{g}/\text{kg}$ in UK Barley from 2015 to 2019 crop years



Diacetoxyscirpenol (DAS) $\mu\text{g}/\text{kg}$ in UK Barley from 2015 to 2019 crop years



Neosolaniol (NEO) $\mu\text{g}/\text{kg}$ in UK Barley from 2015 to 2019 crop years



OTA $\mu\text{g}/\text{kg}$ in UK malt made from barley in crop years 2015 to 2019
 (EU MRL set at 3ppb ($\mu\text{g}/\text{kg}$) for processed cereals in 2003)

